

Packing Tree Degree Sequences

Kristóf Bérczi

Department of Operations Research and MTA-ELTE Egerváry Research Group
Eötvös Loránd University
Pázmány Péter sétány 1/c, 1117 Budapest, Hungary
E-mail: berkri@cs.elte.hu

Zoltán Király

Department of Computer Science and MTA-ELTE Egerváry Research Group
Eötvös Loránd University
Pázmány Péter sétány 1/c, 1117 Budapest, Hungary
E-mail: kiraly@cs.elte.hu

Changshuo Liu

Budapest Semesters in Mathematics
Bethlen Gábor tér 2, 1071 Budapest, Hungary
E-mail: cl20@princeton.edu

István Miklós

Rényi Institute
Reáltanoda u. 13-15, 1053 Budapest, Hungary
E-mail: miklos.istvan@renyi.mta.hu

Keywords: edge-disjoint realizations, packing spanning trees, degree sequences, uniform sampling

Received: October 30, 2018

Special cases of the edge disjoint realizations of two tree degree sequences are considered in this paper. We show that if there is no node which have degree one in both degree sequences, then they always have edge-disjoint caterpillar realizations. By using a probabilistic method, we prove that two tree degree sequences always have edge-disjoint realizations if each vertex is a leaf in at least one of the trees. We also show that the edge-disjoint realization problem is in P for an arbitrary number of tree sequences with the property that each vertex is a non-leaf in at most one of the trees.

On the other hand, we show that the following problem is already NP-complete: given two graphical degree sequences D_1 and D_2 such that D_2 is a tree degree sequence, decide if there exist edge-disjoint realizations of D_1 and D_2 where the realization of D_2 does not need to be a tree.

Finally, we show that efficient approximations for the number of solutions as well as an almost uniform sampler exist for two tree degree sequences if each vertex is a leaf in at least one of the trees.

Povzetek: V članku so obravnavani posebni primeri povezavno-disjunktnih realizaciji dveh zaporedji stopenj dreves. Pokazano je, da, če ni vozlišča stopnje ena v obeh zaporedjih, potem imata zaporedji vedno povezavno- disjunktni gosenični realizaciji. Pokazan je tudi primer, ko je problem NP-poln.

1 Introduction

Packing degree sequences is related to discrete tomography. The central problem of tomography is to reconstruct spatial objects from lower dimensional projections. The discrete 2D version is to reconstruct a colored grid from vertical and horizontal projections. In the simplest version, this problem is to reconstruct the coloring of an $n \times m$ grid with the requirement that each row and column has a specific number of entries for each color. Such colored matrix

can be considered as a factorization of the complete bipartite graph $K_{n,m}$. Indeed, for each color c_i , the 0-1 matrix of size $n \times m$ obtained by replacing c_i by 1 and all other colors by 0 is an adjacency matrix of a simple bipartite graph such that the disjoint union of these simple graphs is $K_{n,m}$. The prescribed number of entries for each color are the degrees of the simple bipartite graphs. Therefore, an equivalent problem is to give a factorization of the complete bipartite graph into subgraphs with prescribed degree sequences.

It is also possible to consider the non-bipartite version of the edge-disjoint realization problem above. Obviously, the sum of the degrees for each vertex must be $n - 1$ when the complete graph K_n is factorized. Therefore, if there are k degree sequences, the last degree sequence is uniquely determined by the first $k - 1$ degree sequences. When $k = 2$, the problem is reduced to the degree sequence problem, and can be solved in polynomial time [3, 4]. Unfortunately the problem becomes NP-complete already for $k = 3$ [1]. However, special cases are polynomially solvable. Such a special case is when one of the degree sequences is almost regular, that is, any two degrees differ by at most 1 [5].

In this paper we consider the case when $k = 3$, and two of the degree sequences are tree degree sequences. It was already known that this case is tractable [6]. Here we present a new result considering special, caterpillar realizations. Another alternative proof is given for a special subclass of pairs of tree degree sequences that can be extended to an arbitrary number of sequences. The size of the solution space and sampling from it is also discussed. As a negative result, we show that deciding the existence of edge-disjoint realizations for two degree sequences D_1 and D_2 is NP-complete even if D_2 is a tree degree sequence (but its realization does not have to be a tree).

2 Preliminaries

In this section we give the definitions and lemmas needed to state the theorems. The central subject of study in this paper is the edge-disjoint realization problem.

Definition 1. A *degree sequence* $D = (d_1, \dots, d_n)$ is a series of non-negative integers. A degree sequence is *graphical* if there is a vertex labeled simple graph $G = (V, E)$ with $V = \{v_1, \dots, v_n\}$ in which the degree of vertex v_i is exactly d_i for $i = 1, \dots, n$. Such graph G is called a *realization* of D . The edge-disjoint realization problem is the following: given a $c \times n$ degree matrix $D = \{(d_{1,1}, \dots, d_{1,n}), (d_{2,1}, \dots, d_{2,n}), \dots, (d_{c,1}, \dots, d_{c,n})\}$, in which each row of the matrix is a degree sequence, decide if there is an ensemble of edge-disjoint realizations of the degree sequences. Such a set of edge-disjoint graphs is called a *realization* of the degree matrix. Given two degree sequences $D = (d_1, \dots, d_n)$ and $F = (f_1, \dots, f_n)$, their *sum* is defined as $D + F = (d_1 + f_1, \dots, d_n + f_n)$.

For sake of completeness, we define tree degree sequences, path sequences and caterpillars.

Definition 2. Let $D = (d_1, \dots, d_n)$ be a degree sequence. Then D is called a *tree sequence* if $\sum_{i=1}^n d_i = 2n - 2$ and each degree is positive. If all of the degrees are 2 except two of them which are 1, then D is called a *path sequence*. A tree is called a *caterpillar* if its non-leaf vertices span a path.

We will use the following complexity classes later on.

Definition 3. A decision problem is in NP if a non-deterministic Turing Machine can solve it in polynomial time. An equivalent definition is that a witness proving the “yes” answer to the question can be verified in polynomial time. A counting problem is in #P if it asks for the number of witnesses of a problem in NP. A counting problem in #P is in FP if there is a polynomial running time algorithm which gives the solution. It is #P-complete if any problem in #P can be reduced to it by a polynomial-time counting reduction.

Definition 4. A counting problem in #P is in FPRAS (Fully Polynomial Randomized Approximation Scheme) if there exists a randomized algorithm such that for any problem instance x and $\epsilon, \delta > 0$, it generates an approximation \hat{f} for the solution f , satisfying

$$P \left(\frac{f}{1 + \epsilon} \leq \hat{f} \leq (1 + \epsilon) \cdot f \right) \geq 1 - \delta,$$

and the algorithm has a time complexity bounded by a polynomial of $|x|$, $1/\epsilon$ and $-\log(\delta)$.

The total variational distance $d_{TV}(p, \pi)$ between two discrete distributions P and π over the set X is defined as

$$d_{TV}(p, \pi) := \frac{1}{2} \sum_{x \in X} |p(x) - \pi(x)|$$

Definition 5. A counting problem in #P is in FPAUS (Fully Polynomial Almost Uniform Sampler) if there exists a randomized algorithm such that for any instance x and $\epsilon > 0$, it generates a random element of the solution space following a distribution P satisfying

$$d_{TV}(p, U) \leq \epsilon,$$

where U is the uniform distribution over the solution space, and the algorithm has a time complexity bounded by a polynomial of $|x|$ and $-\log(\epsilon)$.

The following technical lemma will be used later for constructing edge-disjoint caterpillar realizations.

Lemma 1. For $n \geq 4$, there are two edge-disjoint Hamiltonian paths in the complete graph K_n whose ends are pairwise different.

Proof. Let $V = \{v_1, \dots, v_n\}$, and let the first Hamiltonian path be $v_1, v_2, v_3, \dots, v_n$. We are going to show by induction that there is a second Hamiltonian path H starting at v_2 , ending at v_3 and using no edge between consecutive integers. For $n = 4$, the path $H = v_2, v_4, v_1, v_3$ does the job. Suppose that $n > 4$ and that we have a path H' on vertices v_1, \dots, v_{n-1} between v_2 and v_3 . Since the path has at least three edges, there is an edge $v_i v_j$ for which $i, j < n - 1$. Replace this edge by edges $v_i v_n$ and $v_n v_j$ for getting the desired path H . \square

3 Packing trees

First we consider the problem of edge-disjoint realization of two tree degree sequences without common leaves.

Theorem 2. *Let $D = (d_1, \dots, d_n)$ and $F = (f_1, \dots, f_n)$ be two tree degree sequences such that $\min_i \{d_i + f_i\} \geq 3$. Then D and F have edge-disjoint caterpillar realizations.*

Proof. The proof goes by induction on n . Observe that the smallest possible n is 4 to accommodate at least $4 = 2 \cdot 2$ leaves (note that each tree has at least two leaves). For $n = 4$, the only possible pair of degree sequences is $(2, 2, 1, 1)$ and $(1, 1, 2, 2)$. By Lemma 1, these sequences have edge-disjoint realizations.

If $n > 4$ and both D and F are path sequences, then there exists edge-disjoint Hamiltonian paths, according to Lemma 1.

So we may suppose that at least one of the degree sequences is not a path sequence. As the sum of the degrees in $D + F$ is $4n - 4$, there are at least four indices where $d_j + f_j = 3$. It is not difficult to see that we can select indices i and j such that, possibly after interchanging the roles of D and F , we have $d_i \geq 3$, $d_j = 1$ and $f_j = 2$.

Modify D and F by removing d_j and f_j and decreasing d_i by 1. This modified D' and F' are tree degree sequences without common leaves on $n - 1$ vertices, therefore, by induction, D' and F' have edge-disjoint caterpillar realizations T'_1 and T'_2 , respectively. Modify T'_1 and T'_2 as follows. Add back vertex v_j and connect it to vertex v_i in T'_1 . The tree T_1 thus arising is a realization of D . Take a path P in T'_2 containing all non-leaf vertices and two leaves. Observe that P has at least 3 edges as otherwise F has $n - 2$ leaves, so D has only two, contradicting to $d_i \geq 3$. Hence P has an edge $v_k v_\ell$ such that $k \neq i$ and $\ell \neq i$. For constructing T_2 , replace edge $v_k v_\ell$ of T'_2 by two edges, $v_k v_j$ and $v_j v_\ell$. The tree T_2 thus obtained is a caterpillar, edge-disjoint from T_1 and is a realization of F . \square

The theorem implicitly states that if two tree degree sequences do not share common leaves then their sum is graphical. If the two trees have common leaves, their sum is not necessarily graphical as shown by the example $D = (2, 1, 1)$ and $F = (2, 1, 1)$. Observe that the largest degree in $D + F$ is 4, and there are only 3 vertices.

On the other hand, if the sum of the two sequences happens to be graphical, then they also have edge-disjoint realizations, as was shown by Kundu in [6].

Theorem 3 ([6]). *Let $D = (d_1, \dots, d_n)$ and $F = (f_1, \dots, f_n)$ be two tree degree sequences. Then there exist edge-disjoint tree realizations of D and F if and only if $D + F$ is graphical.*

However, there are tree degree sequences that have edge-disjoint tree realizations but do not have edge-disjoint caterpillar realizations. For example, consider the tree degree sequences

$$D = (5, 2, 2, 2, 2, 2, 1, 1, 1, 1, 1)$$




Figure 1: Edge-disjoint realization of two degree sequences, both of them are $(5, 2, 2, 2, 2, 2, 1, 1, 1, 1, 1)$

and

$$F = (5, 2, 2, 2, 2, 2, 1, 1, 1, 1, 1)$$

According to Theorem 3, these sequences have edge-disjoint realizations as their sum is graphical (see Fig. 1). We claim that they do not have edge-disjoint caterpillar realizations. To see this, observe that in any caterpillar realization, the degree 5 vertices must be connected to at least 3 leaves. However, there are only 5 vertices that are leaves in any of the trees, showing that any pair of caterpillar realizations will share at least one edge.

Theorem 2 considered the case when the leaf vertices of the degree sequences do not coincide. Now we turn to the opposite end, namely when each vertex is a leaf in at least one of the sequences.

Theorem 4. *Let $D = (d_1, \dots, d_n)$ and $F = (f_1, \dots, f_n)$ be tree degree sequences such that $\min(d_i, f_i) = 1$ for all i . Let T_1 and T_2 be random realizations of D and F uniformly distributed. Then the expected number of common edges of T_1 and T_2 is 1.*

Proof. The proof is based on the following lemma.

Lemma 5. *Let T be a random realization of the tree degree sequence $D = (d_1, \dots, d_n)$ (where $n \geq 3$). Then the probability of having an edge between v_i and v_j is*

$$\frac{d_i + d_j - 2}{n - 2}$$

Proof. It is well known that the number of trees with a given degree sequence is

$$\frac{(n - 2)!}{\prod_{k=1}^n (d_k - 1)!} \tag{1}$$

Let T' denote those trees in which v_i and v_j are adjacent. Let f be a mapping from T' to the set of trees with degree sequence

$$(d_1, \dots, d_{i-1}, d_{i+1}, \dots, d_{j-1}, d_{j+1}, \dots, d_n, d_i + d_j - 2)$$

obtained by joining v_i and v_j to a common vertex. The function f is surjective, and each tree is an image $\binom{d_i+d_j-2}{d_i-1}$ times. Therefore the number of trees in which v_i is adjacent to v_j is

$$\frac{(n-3)!}{(d_i+d_j-3)! \prod_{k \neq i,j} (d_k-1)!} \frac{(d_i+d_j-2)!}{(d_i-1)!(d_j-1)!} = \frac{(d_i+d_j-2)(n-3)!}{\prod_{k=1}^n (d_k-1)!}. \tag{2}$$

The probability that v_i is adjacent to v_j is the ratio of (2) and (1), which is indeed

$$\frac{d_i + d_j - 2}{n - 2},$$

thus concluding the proof of the lemma. \square

Now we turn to the proof of the theorem. Let D and F be degree sequences satisfying that each vertex is a leaf in at least one of the trees. Define

$$\begin{aligned} \mathcal{A} &:= \{i \mid d_i > 1 \wedge f_i = 1\}, \text{ and} \\ \mathcal{B} &:= \{i \mid d_i = 1 \wedge f_i > 1\}. \end{aligned}$$

Note that there might be parallel edges in the two trees only between these two sets. The expected number of parallel edges is then

$$\begin{aligned} \sum_{i \in \mathcal{A}} \sum_{j \in \mathcal{B}} \frac{(d_i-1)(f_j-1)}{(n-2)^2} &= \sum_{i \in \mathcal{A}} \frac{d_i-1}{n-2} \sum_{j \in \mathcal{B}} \frac{f_j-1}{n-2} = \\ \sum_{i=1}^n \frac{d_i-1}{n-2} \sum_{j=1}^n \frac{f_j-1}{n-2} &= 1, \end{aligned}$$

since $d_i = 1$ for all $i \in \bar{\mathcal{A}}$, $f_j = 1$ for all $j \in \bar{\mathcal{B}}$, and the sum of the degrees decreased by 1 is $n - 2$ for any tree degree sequence. This finishes the proof of the theorem. \square

Theorem 4 implies a characterization of realizability for a subclass of tree degree sequences.

Corollary 6. *Let $D = (d_1, \dots, d_n)$ and $F = (f_1, \dots, f_n)$ be tree degree sequences such that each vertex is a leaf in at least one of them. Then D and F have edge-disjoint tree realizations if and only if $d_i < n - 1$ and $f_i < n - 1$ for all i .*

Proof. If $\max_i \{d_i\} = n - 1$ or $\max_i \{f_i\} = n - 1$ then $D + F$ is not graphical. On the other hand, if none of the trees is a star, then there are four distinct indices i_1, i_2, j_1 and j_2 such that $i_1, i_2 \in \mathcal{A}$ and $j_1, j_2 \in \mathcal{B}$. Then there exists a pair of trees T_1 and T_2 such that both trees contain edges (v_{i_1}, v_{j_1}) and (v_{i_2}, v_{j_2}) and T_1 realizes D while T_2 realizes F . Indeed, the degree 1 vertices can be connected to any of the non-leaf vertices. This means that the two sequences have realizations having at least 2 common edges, which is above the expected value. Hence there must also exist a realization with less common edges than the expected number, which is 1. That is, there exists an edge-disjoint realization of the two sequences, thus concluding the proof. \square

This theorem will be useful also at generating random realizations, see the next section.

Similar theorem holds for more tree sequences as well. Let us fix again $V = \{v_1, \dots, v_n\}$. We need the following technical preliminary lemma.

Lemma 7. *Let $D = (d_1, \dots, d_n)$ be a tree degree sequence, $n > m > 2$ and $U = \{v_i \mid d_i > 1\}$. Suppose V_1, \dots, V_{m-1} are pairwise disjoint sets in $L = V \setminus U$. Suppose further that $|U| > 1, |V_1| > 1, \dots, |V_{m-1}| > 1$ and $d_i \leq n - m$ for all i . Then there is a tree T realizing D such that its restriction to $U \cup V_j$ is a non-star tree for all j .*

Proof. For any tree realization T , its restriction to $U \cup V_j$ is a tree because outside U there are only leaves. In the case $|U| \geq 4$ we claim that there is a tree realization T such that its restriction to U is not a star. Indeed, if T restricted to U is a star centered at $u \in U$, then, by the degree bound, there is a leaf $w \in L$ not connected to u . Let $u_1 \in U$ denote the unique neighbor of w in the tree, and let u_2 be a third vertex of U . Replacing edges uu_2 and u_1w by edges u_1u_2 and uw gives another tree realization T whose restriction to U is not a star.

For the case $|U| = 2$, let $U = \{v_i, v_j\}$. Connect first v_i to v_j . Now $d_i + d_j = n$, so $d_i \geq m$ and $d_j \geq m$. For each $k \leq m - 1$, connect one vertex of V_k to v_i and another one to v_j . The remaining leaves in L can be distributed easily: connect any $d_i - m$ of them to v_i and the remainder to v_j , giving the aimed tree realization.

Finally suppose $U = \{v_i, v_j, v_k\}$. We may also suppose that $2 \leq d_i \leq d_j \leq d_k$. Connect first v_i and v_j to v_k . It is easy to connect vertices of L to U in such a way that for each $\ell \leq m - 1$, at least one vertex of V_ℓ is connected to either v_j or v_k . \square

Theorem 8. *Let D_1, D_2, \dots, D_m be tree degree sequences with $D_i = d_{i,1}, d_{i,2}, \dots, d_{i,n}$ such that each vertex is a leaf in all except at most one of them. Then D_1, D_2, \dots, D_m have edge-disjoint realizations if and only if $\max_{i,j} \{d_{i,j}\} \leq n - m$.*

Proof. Necessity is clear as $D_1 + D_2 + \dots + D_m$ is not graphical if $\max_{i,j} \{d_{i,j}\} > n - m$.

The statement is trivial when $m = 1$. If $m = 2$ then it is equivalent to Corollary 6, so we may suppose $m > 2$.

We give a constructive proof for the other direction. First a trial solution is built which might contain parallel edges, then these parallel edges are eliminated to get an edge-disjoint realization.

Let V_i denote the subset of vertices on which the degrees in D_i are larger than 1. Note that $\{V_1, V_2, \dots, V_m\}$ forms a subpartition of V and $|V_i| \geq 2$ for each $i = 1, \dots, m$. For a degree sequence D_i , construct a trial tree \tilde{T}_i by using Lemma 7, which ensures that the subtree on vertices $V_i \cup V_k$ is a non-star tree for any $k \neq i$.

From the trial solution, which might contain several parallel edges, a final solution is built in the following way. While there exists a pair of indexes (i, k) such that there

is one or more parallel edges between V_i and V_k , do the following. Let $\tilde{T}_{i,k}$ denote the subtree of the tree \tilde{T}_i on vertices $V_i \cup V_k$ and let $\tilde{D}_{i,k}$ denote its degree sequence. By Corollary 6, $\tilde{D}_{i,k}$ and $\tilde{D}_{k,i}$ have edge-disjoint tree realizations. Replace $\tilde{T}_{i,k}$ and $\tilde{T}_{k,i}$ by such realizations. This removes all parallel edges between V_i and V_k because \tilde{T}_j has no edge between these sets if $j \neq i, j \neq k$. \square

4 Counting and sampling realizations

Since typically there are more than one realizations when a realization exists, and typically the number of realizations might grow exponentially, it is also a computational challenge to estimate their number and/or sample almost uniformly a solution. Here we prove the following theorem.

Theorem 9. *Let $D = (d_1, \dots, d_n)$ and $F = (f_1, \dots, f_n)$ be two tree degree sequences such that each vertex is a leaf in at least one of the trees. Furthermore, assume that none of the trees is a star. Then there is an FPRAS for estimating the number of disjoint realizations and there is an FPAUS for almost uniformly sampling realizations.*

Proof. This theorem is based on Theorem 4. As we discussed, there are random trees with at least two parallel edges. The number of pair of trees containing parallel edges (v_{i_1}, v_{j_1}) and (v_{i_2}, v_{j_2}) such that $d_{i_1}, d_{i_2} > 1$ and $f_{j_1}, f_{j_2} > 1$ is

$$\frac{(n-4)!}{(d_{i_1}-2)!(d_{i_2}-2)! \prod_{k \neq i_1, i_2} (d_k-1)!} \cdot \frac{(n-4)!}{(d_{j_1}-2)!(d_{j_2}-2)! \prod_{k \neq j_1, j_2} (f_k-1)!} \quad (3)$$

Therefore, at least the same number of pair of trees have no parallel edges (that is, are edge-disjoint realizations of the degree sequences) to get the expectation 1 for the number of parallel edges. Therefore, the probability that two random trees will be edge-disjoint is at least

$$\frac{(d_{i_1}-1)(d_{i_2}-1)(f_{j_1}-1)(f_{j_2}-1)}{(n-2)^2(n-3)^2}$$

It follows from basic probabilistic arguments that an FPRAS algorithm can be designed based on this property. Indeed, let ξ be the indicator variable that a random pair of trees are edge-disjoint realizations. Then the number of edge-disjoint realizations is

$$E[\xi] \frac{(n-2)!}{\prod_{k=1}^n (d_k-1)!} \frac{(n-2)!}{\prod_{k=1}^n (f_k-1)!}$$

Furthermore, we know that

$$E[\xi] \geq \frac{(d_{i_1}-1)(d_{i_2}-1)(f_{j_1}-1)(f_{j_2}-1)}{(n-2)^2(n-3)^2}$$

Uniformly distributed random trees with a prescribed degree sequence can be generated in polynomial time based on the fact that the probability that a given leaf is connected to a vertex with degree d_i is

$$\frac{d_i-1}{n-2}$$

A uniformly distributed tree can be generated by randomly selecting a neighbor of a given leaf, then generating a random tree for the remaining degree sequence. Equivalently, the trees with a prescribed degree sequence can be encoded by the Prüfer codes in which the index i appears exactly $d_i - 1$ times. Uniformly generating such Prüfer codes is an elementary computational task.

Therefore, random pair of trees can be generated in polynomial time, and it is easy to check whether or not they are edge-disjoint realizations. Such sampling of random trees provide an unbiased estimation for the expectation of the indicator variable ξ . Indeed, if X_i is 1 if the i^{th} pair of random trees are edge-disjoint and 0 otherwise, then the random variable

$$Y_m := \sum_{i=1}^m X_i$$

follows a binomial distribution with parameter $p = E[\xi]$ and expectation $mE[\xi]$. The tails of the binomial distributions can be bounded by the Chernoff’s inequality:

$$P(Y_m \leq mp(1-\epsilon)) \leq \exp\left(-\frac{1}{2p} \frac{(mp - mp(1-\epsilon))^2}{m}\right)$$

This should be bounded by $\frac{\delta}{2}$ (the other half δ error will go to the other tail)

$$\exp\left(-\frac{1}{2p} \frac{(mp - mp(1-\epsilon))^2}{m}\right) \leq \frac{\delta}{2} \quad (4)$$

Solving Equation 4, we get

$$m \geq \frac{-2 \log\left(\frac{\delta}{2}\right)}{p\epsilon^2}$$

For the upper tail, we can also use the Chernoff’s inequality, just replacing P with $1-p$ and the upper threshold $mp(1+\epsilon)$ with $m - mp(1+\epsilon)$:

$$P(Y_m \geq mp(1+\epsilon)) \leq \exp\left(-\frac{(m(1-p) - (m-mp(1+\epsilon)))^2}{2(1-p)m}\right)$$

Upper bounding this with $\frac{\delta}{2}$ and solving the inequality, we get that

$$m \geq \frac{-2(1-p) \log\left(\frac{\delta}{2}\right)}{p^2\epsilon^2}$$

Since $\frac{1}{p} = O(n^4)$, the necessary number of samples is indeed polynomial with the size of the problem, $\frac{1}{\epsilon}$ and $-\log(\delta)$. Furthermore, one sample can be generated in polynomial time, therefore this algorithm is indeed an FPRAS.

It is also well known that an FPAUS algorithm can be designed in this case. The FPAUS algorithm generate $\frac{-\log(\epsilon)}{p}$ pair of random trees. If any of them is an edge-disjoint realization, then the algorithm returns with it. Otherwise it generates an arbitrary realization and returns with it.

This is indeed an FPAUS algorithm, since any random pair of trees which are edge-disjoint come from sharp the uniform distribution of the solutions. The probability that there will be no edge-disjoint pair of trees in m number of samples is

$$(1 - p)^m.$$

This probability is not larger than ϵ . Indeed,

$$(1 - p)^{\frac{-\log(\epsilon)}{p}} \leq \epsilon,$$

since

$$\frac{-\log(\epsilon)}{p} \log(1 - p) \leq \log(\epsilon)$$

because

$$-\log(1 - p) \geq p.$$

Namely, the algorithm generates realizations from a distribution which is the convex combination $(1 - \epsilon')U + \epsilon'\pi$, where $\epsilon' \leq \epsilon$, U is the uniform distribution and π is an arbitrary distribution. However, the variational distance of this distribution from the uniform one is

$$\begin{aligned} d_{TV}(U, (1 - \epsilon')U + \epsilon'\pi) &= \\ \frac{1}{2} \sum_x |U(x) - ((1 - \epsilon')U(x) + \epsilon'\pi(x))| &= \\ \epsilon' \frac{1}{2} \sum_x |U(x) - \pi(x)| &\leq \epsilon' \leq \epsilon. \end{aligned}$$

Since one sample can be generated in polynomial time, and the total number of samples is polynomial with the size of the problem and $-\log(\epsilon)$, this algorithm is indeed and FPAUS. \square

It remains an open question whether or not similar theorems exist for the case when the tree degree sequences have common high degrees. Also it is open if exact counting of the edge-disjoint solutions is possible in polynomial time, although the natural conjecture is that this counting problem is #P-complete.

5 A complexity result

What can we say when only one of the two degree sequences is a tree degree sequence and the other is arbitrary? Unfortunately, we have a negative result here.

Theorem 10. *It is NP-complete to decide if a tree degree sequence and an arbitrary degree sequence have an edge-disjoint realization (in which the tree degree sequence does not necessarily have to be represented by a tree).*

Proof. By [1], it is NP-complete to decide if two bipartite degree sequences has edge-disjoint realizations. We have the following observations.

Claim 1. *A bipartite degree sequence pair*

$$D = (d_{1,1}, \dots, d_{1,n_1}), (d_{2,1}, \dots, d_{2,n_2})$$

and

$$F = (f_{1,1}, \dots, f_{1,n_1}), (f_{2,1}, \dots, f_{2,n_2})$$

has an edge disjoint realization if and only if the simple degree sequence pair

$$D' = (d_{1,1} + n_1 - 1, \dots, d_{1,n_1} + n_1 - 1, d_{2,1}, \dots, d_{2,n_2})$$

and

$$F' = (f_{1,1}, \dots, f_{1,n_1}, f_{2,1} + n_2 - 1, \dots, f_{2,n_2} + n_2 - 1)$$

has an edge-disjoint realization.

Proof. If an edge-disjoint bipartite realization of D and F is given, then the complete graph on the first vertex class can be added to the first realization and the complete graph on the second vertex class can be added to the second realization to get a (now non-bipartite) realization of D' and F' . On the other hand, it is easy to see that any realization of D' contains K_{n_1} on the first n_1 vertices, and any realization of F' contains K_{n_2} on the last n_2 vertices. Given an edge-disjoint realization of D' and F' , deleting K_{n_1} from D' and K_{n_2} from F' yields an edge-disjoint realization of D and F . \square

Claim 2. *The degree sequence pair $D = (d_1, \dots, d_n)$ and $F = (f_1, \dots, f_n)$ has an edge-disjoint realization if and only if the degree sequence pair $D' = (d_1 + 1, \dots, d_n + 1, n)$ and $F' = (f_1, \dots, f_n, 0)$ has an edge-disjoint realization.*

Proof. Let G_1 and G_2 be an edge-disjoint realization of D and F . Then add a vertex v_{n+1} to G_1 , and connect it to all the other vertices to get a realization of D' . Add an isolated vertex v_{n+1} to G_2 to get a realization of F' . These realizations of D' and F' are edge-disjoint. On the other hand, in any realization of D' , v_{n+1} is connected to all the other vertices. If edge-disjoint realizations of D' and F' are given, delete v_{n+1} from both realizations to get edge-disjoint realizations of D and F . \square

Claim 3. *The degree sequence pair $D = (d_1, \dots, d_n)$ and $F = (f_1, \dots, f_n)$ has an edge-disjoint realization if and only if the degree sequence pair $D' = (d_1, \dots, d_n, 1, 1)$ and $F' = (f_1 + 1, \dots, f_n + 1, n, 0)$ has an edge-disjoint realization.*

Proof. Any edge-disjoint realization G_1 and G_2 of D and F can be extended to an edge-disjoint realization of D' and F' by adding two vertices v_{n+1} and v_{n+2} , and then connecting v_{n+1} to all v_1, \dots, v_n in G_2 and connecting v_{n+1} and v_{n+2} in G_1 . On the other hand, in any edge-disjoint realizations G'_1 and G'_2 of D' and F' , v_{n+1} is connected to all v_1, \dots, v_n in G'_2 , therefore, v_{n+1} must be connected to v_{n+2} in G'_1 . Therefore deleting v_{n+1} and v_{n+2} yields an edge-disjoint realization of D and F . \square

We can use Claim 1 to prove that it is NP-complete to decide if two simple degree sequences have edge-disjoint realizations. Claim 2 shows that it is NP-complete to decide if two degree sequences have edge-disjoint realizations such that one of the degree sequences does not have 0 degrees. Finally, Claim 3 can be used to iteratively transform any D degree sequence (that already does not have a 0 degree) into a tree degree sequence. Indeed, in each step, we add two vertices to D and extend the sum of the degrees only by 2. Therefore in a polynomial number of steps, we get a degree sequence D' in which the sum of the degrees is exactly twice the number of vertices minus 2. Therefore it follows that given any bipartite degree sequences D and F , we can construct in polynomial time two simple degree sequences D' and F' such that D and F have edge-disjoint realizations if and only if D' and F' have edge-disjoint realizations, furthermore, D' is a tree degree sequence. \square

6 Discussion and conclusions

In this paper, we considered packing tree degree sequences. When there are no common leaves, there are always edge-disjoint caterpillar realizations. On the other hand, there might not be edge-disjoint caterpillar realizations when there are common leaves, even if otherwise there are edge-disjoint tree realizations.

When there are no common high degree vertices, there are edge-disjoint tree realizations if and only if none of the degree sequences is a degree sequence of a star. Similar theorem exists for arbitrary number of trees, and it is easy to decide if arbitrary number of tree degree sequences without common high degrees have edge-disjoint realizations.

It is also known [5] that a degree sequence and an almost regular degree sequence have an edge-disjoint realization if and only if their sum is graphical. This raises the natural question if a degree sequence and a tree sequence have edge-disjoint realizations if and only if their sum is graphical. We showed that the answer is no to this question, and actually it is NP-complete to decide if an arbitrary degree sequence and a tree degree sequence have edge-disjoint realizations.

We also showed how to approximately count and sample edge-disjoint tree realizations with prescribed degrees. We proved that this is possible if there are no common high degree vertices. However, when the two degree sequences have common high degree vertices then the problem remains open.

References

- [1] Dürr, C., Guinez, F., Matamala, M.: Reconstructing 3-colored grids from horizontal and vertical projections is NP-hard. *European Symposium on Algorithms*, 776–787 (2009)
- [2] Erdős, P.; Gallai, T. : Graphs with vertices of prescribed degrees (in Hungarian) *Matematikai Lapok*, 11: 264–274. (1960)
- [3] S.L. Hakimi: On the degrees of the vertices of a directed graph. *J. Franklin Institute*, 279(4):290–308. (1965)
- [4] V. Havel: A remark on the existence of finite graphs. (Czech), *Časopis Pěst. Mat.* 80:477–480. (1955)
- [5] Kundu, S.: The k-factor conjecture is true. *Discrete Mathematics*, 6(4):367–376. (1973)
- [6] Kundu, S.: Disjoint Representation of Tree Realizable Sequences. *SIAM Journal on Applied Mathematics*, 26(1):103–107. (1974)

